

**Datasheet for 611-1223****Rabbit IgG Fc Antibody Fluorescein Conjugated Pre-Adsorbed****Overview**

|                      |                                                                                                 |
|----------------------|-------------------------------------------------------------------------------------------------|
| <b>Description:</b>  | Goat Anti-Rabbit IgG Fc Antibody Fluorescein Conjugated (Min X Human Serum Proteins) - 611-1223 |
| <b>Item No.:</b>     | 611-1223                                                                                        |
| <b>Size:</b>         | 2 mg                                                                                            |
| <b>Reactivity:</b>   | Rabbit                                                                                          |
| <b>Host Species:</b> | Goat                                                                                            |

**Product Details**

|                           |                                                                                                                                                                                                                                                                                                                                                                                                 |
|---------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Background:</b>        | Anti-Rabbit IgG F(c) Fluorescein Antibody generated in goat is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc portion of rabbit IgG and often this fragment is removed from immunoglobulins to minimize receptor binding and lower background reactivity. |
| <b>Synonyms:</b>          | Goat anti-Rabbit IgG F(c) Antibody fluorescein Conjugation, Goat anti-Rabbit IgG Fc fragment Antibody fluorescein Conjugation, Goat anti-Rabbit IgG F(c) FITC Conjugated Antibody                                                                                                                                                                                                               |
| <b>Host Species:</b>      | Goat                                                                                                                                                                                                                                                                                                                                                                                            |
| <b>Specificity:</b>       | IgG Fc                                                                                                                                                                                                                                                                                                                                                                                          |
| <b>Conjugate:</b>         | Fluorescein (FITC)                                                                                                                                                                                                                                                                                                                                                                              |
| <b>Clonality:</b>         | Polyclonal                                                                                                                                                                                                                                                                                                                                                                                      |
| <b>Format:</b>            | IgG                                                                                                                                                                                                                                                                                                                                                                                             |
| <b>F/P Ratio:</b>         | 7.23                                                                                                                                                                                                                                                                                                                                                                                            |
| <b>Specific Activity:</b> | 2.75                                                                                                                                                                                                                                                                                                                                                                                            |

**Target Details**

|                    |                          |
|--------------------|--------------------------|
| <b>Reactivity:</b> | Rabbit                   |
| <b>Immunogen:</b>  | Rabbit IgG F(c) fragment |

|                            |                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
|----------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Purity/Specificity:</b> | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Goat Serum, Rabbit IgG, Rabbit IgG F(c) and Rabbit Serum. No reaction was observed against Rabbit IgG F(ab) or Human Serum Proteins. |
|----------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|

## Application Details

|                          |                                                                                                                                                                                                                                                                   |
|--------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Application Note:</b> | This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms. |
| <b>Assay Dilutions:</b>  | All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.                                                                                                                                                                   |
| <b>FC:</b>               | 1:500 - 1:2,500                                                                                                                                                                                                                                                   |
| <b>FLISA:</b>            | 1:10,000 - 1:50,000                                                                                                                                                                                                                                               |
| <b>IF:</b>               | 1:1,000 - 1:5,000                                                                                                                                                                                                                                                 |


## Formulation

|                               |                                                                        |
|-------------------------------|------------------------------------------------------------------------|
| <b>Physical State:</b>        | Lyophilized                                                            |
| <b>Concentration:</b>         | 2 mg/mL by UV absorbance at 280 nm                                     |
| <b>Buffer:</b>                | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2             |
| <b>Preservative:</b>          | 0.01% (w/v) Sodium Azide                                               |
| <b>Stabilizer:</b>            | 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free |
| <b>Reconstitution Volume:</b> | 1.0 mL                                                                 |
| <b>Reconstitution Buffer:</b> | Restore with deionized water (or equivalent)                           |

## Shipping & Handling

|                            |                                                                                                                                                                                                                                                                                                                                                 |
|----------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Shipping Condition:</b> | Ambient                                                                                                                                                                                                                                                                                                                                         |
| <b>Storage Condition:</b>  | Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. |
| <b>Expiration:</b>         | Expiration date is one (1) year from date of receipt.                                                                                                                                                                                                                                                                                           |

## Images



### Western Blot

Western Blot of Anti-Rabbit IgG F(c) (GOAT) Antibody (p/n 611-1103). Lane M: 3  $\mu$ l Molecular Ladder. Lane 1: Rabbit IgG whole molecule (p/n 011-0102). Lane 2: Rabbit IgG F(ab) Fragment (p/n 011-0105). Lane 3: Rabbit IgG F(c) Fragment (p/n 010-0103). Lane 4: Rabbit IgM Whole Molecule (p/n 011-0107). Lane 5: Normal Rabbit Serum (p/n B309). All samples were reduced. Load: 50 ng of IgG, F(ab), IgM and Serum, 100 ng of F(c). Block: MB-070 for 30 min at RT. Primary Antibody: Anti-Rabbit IgG F(c) (GOAT) Antibody (p/n 611-1103) 1:2,000 for 60 min at RT. Secondary antibody: Anti-Goat IgG (DONKEY) Peroxidase Conjugated Antibody (p/n CUST10) 1:40,000 in MB-070 for 30 min at RT. Predicted/Observed Size: 25 and 50 kDa for Rabbit IgG and Serum, 25 kDa for F(c) and F(ab), 70 and 23 kDa for IgM. Rabbit F(c) migrates slightly higher.

## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.