Order online now!

www.rockland.com
tech@rockland.com
+1 484.791.3823

Datasheet for 611-201-122

Rabbit IgG (H&L) Antibody Pre-Adsorbed

Overview

Description:	Guinea Pig Anti-Rabbit IgG (H&L) Antibody (Min X Hu, Gt, Ms Serum Proteins) - 611-201-122
Item No.:	611-201-122
Size:	1 mg
Applications:	ELISA, WB, CUT&RUN
Reactivity:	Rabbit
Host Species:	Guinea Pig

Product Details

Background:	Anti-Rabbit IgG (H&L) generated in guinea pig detects rabbit Immunoglobulin G. Both the Heavy and Light chains of the antibody molecule are present. Representing approximately 75% of serum immunoglobulins, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized and secreted by plasma B cells. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.
Synonyms:	Guinea Pig Anti-Rabbit IgG Antibody, Guinea Pig Anti Rabbit IgG Antibody
Host Species:	Guinea Pig
Specificity:	IgG (H&L)
Clonality:	Polyclonal
Format:	IgG

Target Details

Reactivity:	Rabbit
Immunogen:	Rabbit IgG whole molecule

Purity/Specificity:	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Guinea Pig Serum, Rabbit IgG and Rabbit Serum. No reaction was observed against Goat, Human and Mouse Serum Proteins.

Application Details

Tested Applications:	ELISA, WB
Suggested Applications:	CUT&RUN (Based on references)
Application Note:	Anti-Rabbit IgG antibody has been tested by ELISA and Western Blot and is suitable for use in immunohistochemistry and CUT&RUN. Specific conditions for reactivity should be optimized by the end user.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:15,500 - 1:17,500
IHC:	1:1,000 - 1:5,000
WB:	1:2,000 - 1:10,000

Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1.01 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	None

Shipping & Handling

Shipping Condition:	Wet Ice
Storage Condition:	Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.
Expiration:	Expiration date is one (1) year from date of receipt.

Images



ELISA
ELISA Results of Purified Guinea Pig Anti-Rabbit IgG mx3 Antibody tested against purified Rabbit IgG (Mx Human IgG, Goat IgG, Mouse IgG). Each well was coated in duplicate with 10 µg of Rabbit IgG (p/n 011-0102) [Red Line], Human IgG (p/n 009-0102) [Green Line], Goat IgG (p/n 005-0102) [Blue Line], Mouse IgG (p/n 010-0102) [Purple Line]. The working dilution is 1:17,000. The starting dilution of antibody was 5µg/ml and the X-axis represents the Log10 of a 3-fold dilution. This titration is a 4-parameter curve fit where the IC50 is defined as the titer of the antibody. Assay performed using 3% Fish Gel/PBS Blocking buffer (p/n MB-066), Goat Anti-Guinea Pig mx10 HRP conjugated (p/n 606-103-129) and TMB substrate (p/n TMBE-1000).



Western Blot
Western Blot of Anti-Rabbit IgG (H&L) (GUINEA PIG) Antibody (Min X Hu, Gt, Ms Serum Proteins) (p/n 611-201-122). Lane M: 3 µl Molecular Ladder. Lane 1: Rabbit IgG whole molecule (p/n 011-0102). Lane 2: Rabbit IgG F(ab) Fragment (p/n 011-0105). Lane 3: Rabbit IgG F(c) Fragment (p/n 010-0103). All samples were reduced. Load: 50 ng per lane. Block: MB-070 for 30 min at RT. Primary Antibody: Anti-Rabbit IgG (H&L) (GUINEA PIG) Antibody (Min X Hu, Gt, Ms Serum Proteins) (p/n 611-201-122) 1:500 for 60 min at RT. Secondary antibody: Anti-Guinea Pig IgG (GOAT) Peroxidase Conjugated Antibody (p/n 606-103-129) 1:40,000 in MB-070 for 30 min at RT. Predicted/Observed Size: 25 and 50 kDa for Rabbit IgG and Serum, 25 kDa for F(c) and F(ab). Rabbit F(c) migrates slightly higher.

References

- Kujirai T et al. Structural insights into how DEK nucleosome binding facilitates H3K27 trimethylation in chromatin. *Nat Struct Mol Biol.* (2025)
- Kong Y et al. Targeting the SP/KLF Transcriptional Regulatory Network Synergizes with HDAC Inhibition to Impede Progression of H3K27M Diffuse Intrinsic Pontine Glioma. *Cancer Res.* (2025)
- Matsumoto M et al. Role of PRC2 in the stochastic expression of Aire target genes and development of mimetic cells in the thymus. *J Exp Med.* (2025)
- Choksi SP et al. An alternative cell cycle coordinates multiciliated cell differentiation. *Nature.* (2024)
- Xue Y et al. Unraveling the key role of chromatin structure in cancer development through epigenetic landscape characterization of oral cancer. *Mol Cancer.* (2024)
- Fang Y et al. AIRE relies on Z-DNA to flag gene targets for thymic T cell tolerization. *Nature.* (2024)
- Iwagawa T et al. The histone H3K36 demethylase Fbxl11 plays pivotal roles in the development of retinal late-born cell types. *Genes Cells.* (2023)
- Michelson DA et al. Thymic epithelial cells co-opt lineage-defining transcription factors to eliminate autoreactive T cells. *Cell.* (2022)
- Zhang Q et al. Chromatin Accessibility Predetermines Odontoblast Terminal Differentiation. *Front Cell Dev Biol.* (2021)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.