

Datasheet for 612-401-D23**Connexin 43 phospho S368 Antibody****Overview**

Description:	Anti-Connexin 43 pS368 (Rabbit) Antibody - 612-401-D23
Item No.:	612-401-D23
Size:	100 µL
Applications:	WB
Reactivity:	Mouse, Rat
Host Species:	Rabbit

Product Details

Background:	Anti-Connexin 43 pS368 Antibody detects pS368 Connexin which is thought to play a key role in development and may also be involved in epilepsy. Connexin 43 antibody recognizes Connexin43 which forms gap-junctional channels and regulates the permeability of these gap junctions to small organic molecules. Permeability of connexin43 is known to be regulated by phosphorylation at Ser368 by protein kinase C. Phosphorylation of Ser368 by PKC induces a conformational change of connexin 43 that results in a decrease in gap junction permeability. Anti-Connexin 43 pS368 Antibody is ideal for investigators involved in Neuroscience and Cell Signaling.
Synonyms:	Gap junction alpha-1 protein, Gap junction 43 kDa heart protein, CX43, Connexin-43
Host Species:	Rabbit
Clonality:	Polyclonal
Format:	IgG

Target Details

Gene Name:	Gja1
Reactivity:	Mouse, Rat
PTM Specificity:	Phosphorylation
Immunogen Type:	Conjugated Peptide

Immunogen:	Anti-Connexin 43 pS368 Antibody was produced in rabbit by repeated immunizations with synthetic phospho-peptide corresponding to amino acid residues surrounding Ser368 conjugated to KLH.
Purity/Specificity:	Anti-Connexin 43 pS368 antibody is directed against rat pS368 Connexin 43. Anti-Connexin 43 pS368 antibody was prepared from monospecific antiserum by immunoaffinity chromatography using phospho peptide coupled to agarose beads followed by solid phase adsorption(s) against non-phospho peptide and non-specific peptide to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum. This antibody is specific for phosphorylated Connexin 43. Minimal reactivity occurs against non-phosphorylated Connexin 43. Reactivity is completely blocked by treatment with lambda-phosphatase. Reactivity against Connexin 43 occurs from rat sources. However, reactivity is also expected against bovine, canine, chicken, guinea pig, human, mouse and zebra fish sources.
Relevant Links:	• UniProtKB - P08050 • GeneID - 24392 • NCBI - AAA41000.1

Application Details

Tested Applications:	WB
Application Note:	Anti-Connexin 43 pS368 Antibody has been tested in western blot and is suitable for use in ELISA. Specific conditions for reactivity should be optimized by the end user. Expect a band at approximately 43 kDa in size corresponding to connexin 43 protein phosphorylated at Ser368 in the appropriate cell lysate or extract.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:10,000
WB:	1:1000

Formulation


Physical State:	Liquid
Buffer:	0.01 M HEPES, 0.15 M Sodium Chloride, pH 7.5
Stabilizer:	0.1 mg/ml Bovine Serum Albumin (BSA) - IgG and Protease free, 50% (v/v) Glycerol

Shipping & Handling

Shipping Condition:	Dry Ice
----------------------------	---------

Storage Condition:	Store vial at -20° C prior to opening. This product is stable at 4° C as an undiluted liquid. For extended storage, aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Images



Western Blot

Western Blot of Anti-Connexin 43 pS368 (Rabbit) Antibody. Lane 1: rat hippocampal lysate. Lane 2: rat hippocampal lysate incubated in λ-Ptase (1200 units for 30 min). Load: 10 µg per lane. Primary antibody: Connexin 43 pS368 antibody at 1:400 for overnight at 4°C. Secondary antibody: IRDye800™ rabbit secondary antibody at 1:10,000 for 45 min at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: ~43kDa/~43kDa for connexin43 phosphorylated at Ser368. Other band(s): none.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.