ROCKLAND

Order online now!

www.rockland.com
tech@rockland.com
+1 484.791.3823

Datasheet for 702-406-002

F(ab')2 Cat IgG (H&L) Antibody Biotin Conjugated

Overview

Description:	Rabbit F(ab')2 Anti-Cat IgG (H&L) Antibody Biotin Conjugated - 702-406-002
Item No.:	702-406-002
Size:	500 µg
Reactivity:	Cat
Host Species:	Rabbit

Product Details

Background:	This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms. F(ab')2 Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and experimental applications. F(ab)2 fragments penetrate tissue samples and show better antigen recognition and signal generation in IHC. F(ab)2 fragments lack the Fc region and therefore do not bind Fc receptors which effectively lowers background staining. F(ab')2 Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays.
Synonyms:	Rabbit F(ab')2 Anti-Cat biotin Conjugated Antibody, Rabbit Fab2 Anti-Cat Antibody biotin Conjugation, Rabbit Fab'2 Anti-Cat Antibody biotin Conjugation
Host Species:	Rabbit
Specificity:	IgG (H&L)
Conjugate:	Biotin
Clonality:	Polyclonal
Format:	IgG F(ab')2

Target Details

Reactivity:	Cat
Immunogen:	Cat IgG whole molecule

Purity/Specificity:	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Cat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Biotin, anti-Rabbit Serum, Cat IgG and Cat Serum. No reaction was observed against anti-Pepsin or anti-Rabbit IgG F(c).

Application Details

Application Note:	Secondary antibody reagents are ideal for ELISA, western blotting, Immunohistochemistry, Fluorescence Microscopy, Flow Cytometry as well as other antibody detection methods.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:450,000
IHC:	1:1,000 - 1:5,000
WB:	1:10,000 - 1:40,000

Formulation

Physical State:	Lyophilized
Concentration:	1.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Reconstitution Volume:	500 µL
Reconstitution Buffer:	Restore with deionized water (or equivalent)

Shipping & Handling

Shipping Condition:	Ambient
Storage Condition:	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

www.rockland.com
tech@rockland.com
+1 484.791.3823

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.