

Datasheet for 703-402-002

F(ab')₂ Chicken IgG (H&L) Antibody Fluorescein Conjugated**Overview**

| | |
|----------------------|---|
| Description: | Rabbit F(ab') ₂ Anti-Chicken IgG (H&L) Antibody Fluorescein Conjugated - 703-402-002 |
| Item No.: | 703-402-002 |
| Size: | 500 µL |
| Reactivity: | Chicken |
| Host Species: | Rabbit |

Product Details

| | |
|----------------------|---|
| Background: | F(ab') ₂ Anti-Chicken IgG Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab') ₂ fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and experimental applications. F(ab') ₂ fragments penetrate tissue samples and show better antigen recognition and signal generation in IHC. F(ab') ₂ fragments lack the Fc region and therefore do not bind Fc receptors which effectively lowers background staining. F(ab') ₂ Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays. |
| Synonyms: | Rabbit F(ab') ₂ Anti-Chicken IgG FITC Conjugated Antibody, Rabbit F(ab') ₂ Anti-Chicken IgY FITC, Rabbit Fab2 Anti Chicken Fluorescein Conjugated Antibody, Rabbit Fab'2 Anti-Chicken Antibody Fluorescein Conjugation |
| Host Species: | Rabbit |
| Specificity: | IgG (H&L) |
| Conjugate: | Fluorescein (FITC) |
| Clonality: | Polyclonal |
| Format: | IgG F(ab') ₂ |
| F/P Ratio: | 2.1 |

Target Details

| | |
|--------------------|----------------------------|
| Reactivity: | Chicken |
| Immunogen: | Chicken IgG whole molecule |

Purity/Specificity: This product was prepared from monospecific antiserum by immunoaffinity chromatography using Chicken IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein, anti-Rabbit Serum, Chicken IgG and Chicken Serum. No reaction was observed against anti-Pepsin or anti-Rabbit IgG F(c).

Application Details

Application Note: Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.

Assay Dilutions: All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

FC: 1:500-1:2,500

FLISA: 1:10,000 - 1:50,000

IF: 1:1,000 - 1:5,000

Formulation

Physical State: Lyophilized

Concentration: 1.0 mg/mL by UV absorbance at 280 nm

Buffer: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Preservative: 0.01% (w/v) Sodium Azide

Stabilizer: 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free

Reconstitution Volume: 500 μ L

Reconstitution Buffer: Restore with deionized water (or equivalent)

Shipping & Handling

Shipping Condition: Ambient

Storage Condition: Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Expiration: Expiration date is one (1) year from date of receipt.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.