

Datasheet for 709-1116**F(ab')₂ Human IgG Antibody Pre-Adsorbed****Overview**

Description:	Goat F(ab') ₂ Anti-Human IgG (H&L) Antibody (Min X Bv Hs Ms & Rt Serum Proteins) - 709-1116
Item No.:	709-1116
Size:	1 mg
Applications:	Dot Blot, ELISA
Reactivity:	Human
Host Species:	Goat

Product Details

Background:	F(ab') ₂ Anti-Human IgG (H&L) Antibody generated in goat detects immunoglobulin g from human, both heavy and light chains of the antibody molecule are present. Each IgG has two antigen binding sites. Representing approximately 75% of serum immunoglobulins in humans, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized and secreted by plasma B cells. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition. F(ab') ₂ Antibody is ideal for investigators who routinely perform flow cytometry, immunohistochemistry or IHC and other immunoassays. F(ab') ₂ Anti-Human IgG (H&L) Antibody is ideal for investigators in Immunology, Cancer, and Microbiology research.
Synonyms:	goat F(ab') ₂ anti-Human IgG Antibody, Gt-a-Human F(ab') ₂ IgG (H&L), Human F(ab') ₂ IgG (H&L) Antibody in goat, Goat IgG (H&L) Secondary Antibody.
Host Species:	Goat
Specificity:	IgG (H&L)
Clonality:	Polyclonal
Format:	IgG F(ab') ₂

Target Details

Reactivity:	Human
--------------------	-------

Immunogen Type:	Native Protein
Immunogen:	Anti-Human IgG was produced by repeated immunization with human IgG whole molecule in goat.
Purity/Specificity:	F(ab') ₂ Anti-Human IgG (H&L) Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Human IgG and Human Serum. No reaction was observed against anti-Pepsin, anti-Goat IgG F(c), or Bovine, Horse, Mouse and Rat Serum Proteins.

Application Details

Tested Applications:	Dot Blot, ELISA
Application Note:	F(ab') ₂ Anti-Human IgG (H&L) Antibody has been tested by ELISA and dot blot and is suitable for ELISA, immunoblotting, and immunohistochemistry as well as other peroxidase-antibody based enzymatic assays requiring lot-to-lot consistency.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:42,000
IHC:	1:500 - 1:2,000
WB:	1:1,000 - 1:5,000

Formulation

Physical State:	Liquid (sterile filtered)
Concentration:	1.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	None

Shipping & Handling

Shipping Condition:	Wet Ice
Storage Condition:	Store vial at 4° C prior to opening. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing.

Expiration: Expiration date is one (1) year from date of receipt.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.