Order online now!

www.rockland.com
tech@rockland.com
+1 484.791.3823

Datasheet for 809-1002

# Fab Human IgG (H&L) Antibody Rhodamine Conjugated

## Overview

| | |
|---|---|
| **Description:** | Goat Fab Anti-Human IgG (H&L) Antibody Rhodamine Conjugated - 809-1002 |
| **Item No.:** | 809-1002 |
| **Size:** | 1 mg |
| **Reactivity:** | Human |
| **Host Species:** | Goat |

## Product Details

| | |
|---|---|
| **Background:** | Fab Anti-Human IgG (H&L) Rhodamine Antibody generated in goat detects immunoglobulin g from human, both heavy and light chains of the antibody molecule are present. Each IgG has two antigen binding sites. Representing approximately 75% of serum immunoglobulins in humans, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized and secreted by plasma B cells. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition. |
| **Synonyms:** | Goat Fab Anti-Human IgG Antibody Rhodamine Conjugation, Goat Fab Anti-Human IgG TRITC Conjugated Antibody |
| **Host Species:** | Goat |
| **Specificity:** | IgG (H&L) |
| **Conjugate:** | Rhodamine (TRITC) |
| **Clonality:** | Polyclonal |
| **Format:** | IgG Fab |
| **F/P Ratio:** | 1.2 |

## Target Details

| | |
|---|---|
| **Reactivity:** | Human |
| **Immunogen:** | Human IgG whole molecule |

| | |
|---|---|
| **Purity/Specificity:** | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, papain digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Fluorescein and anti-Goat Serum. No reaction was observed against anti-Papain or anti-Goat IgG F(c). |

## Application Details

| | |
|---|---|
| **Application Note:** | Fab Anti-Human IgG (H&L) Rhodamine Antibody is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms. Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. |
| **Assay Dilutions:** | All assays should be optimized by the user. Recommended dilutions (if any) may be listed below. |
| **FC:** | 1:500 - 1:2,500 |
| **FLISA:** | 1:10,000 - 1:50,000 |
| **IF:** | 1:1,000 - 1:5,000 |

## Formulation

| | |
|---|---|
| **Physical State:** | Lyophilized |
| **Concentration:** | 1.0 mg/mL by UV absorbance at 280 nm |
| **Buffer:** | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| **Preservative:** | 0.01% (w/v) Sodium Azide |
| **Stabilizer:** | 10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free |
| **Reconstitution Volume:** | 1.0 mL |
| **Reconstitution Buffer:** | Restore with deionized water (or equivalent) |

## Shipping & Handling

| | |
|---|---|
| **Shipping Condition:** | Ambient |

| | |
|---|---|
| **Storage Condition:** | Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use. |
| **Expiration:** | Expiration date is one (1) year from date of receipt. |

## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.