

## Datasheet for 809-4102

**Fab Human IgG (H&L) Antibody****Overview**

<b>Description:</b>	Rabbit Fab Anti-Human IgG (H&L) Antibody - 809-4102
<b>Item No.:</b>	809-4102
<b>Size:</b>	500 µg
<b>Applications:</b>	IF
<b>Reactivity:</b>	Human
<b>Host Species:</b>	Rabbit

**Product Details**

<b>Background:</b>	Fab Anti-Human IgG (H&L) Antibody generated in rabbit detects immunoglobulin g from human, both heavy and light chains of the antibody molecule are present. Each IgG has two antigen binding sites. Representing approximately 75% of serum immunoglobulins in humans, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized and secreted by plasma B cells. Secondary Antibodies are available in a variety of formats and conjugate types. When choosing a secondary antibody product, consideration must be given to species and immunoglobulin specificity, conjugate type, fragment and chain specificity, level of cross-reactivity, and host-species source and fragment composition.
<b>Synonyms:</b>	Rabbit Fab Anti-Human IgG Antibody, Rabbit Fab Fragment Anti-Human IgG Antibody
<b>Host Species:</b>	Rabbit
<b>Specificity:</b>	IgG (H&L)
<b>Clonality:</b>	Polyclonal
<b>Format:</b>	IgG Fab

**Target Details**

<b>Reactivity:</b>	Human
<b>Immunogen:</b>	Human IgG whole molecule

**Purity/Specificity:** This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, papain digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum. No reaction was observed against anti-Papain or anti-Rabbit IgG F(c).

## Application Details

**Suggested Applications:** IF (Based on references)

**Application Note:** Fab Anti-Human IgG (H&L) Antibody is suitable for highly specific immunological methods requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.

**Assay Dilutions:** All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

**ELISA:** 1:30,000

**IHC:** 1:1,000 - 1:5,000

**WB:** 1:2,000 - 1:10,000

## Formulation

**Physical State:** Liquid (sterile filtered)

**Concentration:** 1.0 mg/mL by UV absorbance at 280 nm

**Buffer:** 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

**Preservative:** 0.01% (w/v) Sodium Azide

**Stabilizer:** None


## Shipping & Handling

**Shipping Condition:** Wet Ice

**Storage Condition:** Store vial at 4° C prior to opening. This product is stable at 4° C as an undiluted liquid. Dilute only prior to immediate use.

**Expiration:** Expiration date is one (1) year from date of receipt.

## Images



### Immunofluorescence Microscopy

Fluorescence microscopy using Rabbit Fab Anti-Human IgG. Analysis of human IgGs displayed on the surface of OCMS™ hybridomas. (a) 8C5 OCMS™ (top row) and B5-6T OCMS™ (bottom row) cells were incubated overnight with the RAH (AB14), followed by biotinylated rabies GP and Alexa Fluor 488® streptavidin (green) (AB3), and visualized by fluorescence microscopy. Images are shown with DAPI (blue) (right) or without DAPI (left). Scale bar = 5 μm. (b) The 4G4 OCMS™ (top row) and 8C5 OCMS™ (bottom row) hybridomas were incubated overnight with a rabbit Fab anti-human IgG (AB17), then for an hour with additional rabbit Fab anti-human IgG and Sabin type III PV (left column) or GP (right column), followed by an Alexa Fluor® 488 anti-human IgG (green) (AB2). Nuclei were stained with DAPI (blue). Scale bar = 10 μm. (c) In this mixing experiment, non-secretor OCMS™ cell lines were labeled with CFSE (green) and mixed in a 10:1 ratio with IgG-secreting OCMS™ hybridomas. Cells were incubated overnight with rabbit F(ab')<sub>2</sub> anti-human IgG (AB15) then washed and incubated with biotinylated Protein A (AB12) and APC streptavidin (red) (AB4). (Top row) 8C5 OCMS™ vs. B5-6T OCMS™ CFSE. (Bottom row) 4G4 OCMS™ vs. LCX OCMS™ CFSE. With DAPI (blue) (right column) or without (left column). Scale bar = 5 μm. Figure 2. PMID: 30794061.

## References

- Puligedda RD et al. Capture and display of antibodies secreted by hybridoma cells enables fluorescent on-cell screening. *MABs*. (2019)

## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.