

Datasheet for KCAE75

AKT3 Western Chemiluminescent Blotting Kit**Overview**

Description:	AKT3 Western Blot Kit - KCAE75
Item No.:	KCAE75
Size:	1 Kit
Applications:	IF, WB
Reactivity:	Human
Host Species:	Mouse

Product Details

Background:	Rockland Immunochemicals' Chemiluminescent Western Blot Kit for AKT3 combines all of the necessary reagents with a rapid proven protocol and the extremely high signal detection of our luminol chemiluminescent substrate for the detection of native and/or recombinant human AKT3 protein. AKT3 Western Blotting kit allows for the detection of endogenous protein levels of human AKT3 present in cell lysates provided by the user. As a positive control, this kit includes a MDA-MB468 whole cell lysate proven to contain AKT3. After protein separation by SDS-PAGE and transfer, the membrane is probed with Rockland's optimized Anti-AKT3 antibody. Detection of the membrane bound antibody-antigen complex is achieved by the addition of a secondary antibody conjugated to the enzyme horseradish peroxidase. The enzyme reacts with a specialized formulation of luminol, an extremely sensitive, non-radioactive substrate that emits light and allows visualization using X-ray film or other imaging methods, including highly sensitive CCD cameras and imaging systems. AKT3 Western Blotting is ideal for investigators involved in Cell Signaling, Cancer, Neuroscience and Signal Transduction research.
Synonyms:	AKT 3 western blotting, AKT-3, PKB antibody, PKB gamma antibody, PKBGAMMA antibody, PRKBG antibody, Protein kinase Akt 3 antibody, Protein kinase B gamma antibody, RAC-gamma serine/threonine-protein kinase, RAC-PK-gamma, western blotting kit, Chemiluminescent WB, Western Blotting Kit, Chemiluminescent Kit, Peroxidase Kit, Immunoblotting Kit
Host Species:	Mouse
Clonality:	Monoclonal
Clone ID:	9A5.H9.G7
Format:	IgG1
Detection Kit Type:	Chemiluminescent Western Blot Kit

Target Details

Gene Name:	AKT3
Reactivity:	Human
Purity/Specificity:	The kit is designed to detect both unphosphorylated and phosphorylated forms of the protein. Cross reactivity with AKT3 from other species has not been determined, however, the sequence of the immunogen shows 100% identity to human, mouse, and rat, therefore, cross reactivity is expected. Cross-reactivity with AKT1 and AKT2 has not been determined. This kit contains sufficient substrate for up to 5 mini blots at 7.5 x 8 cm ² (1,800 cm ²) and is stable for at least 1 year when stored as indicated.
Relevant Links:	<ul style="list-style-type: none">• GeneID - 10000• UniProtKB - Q9Y243• NCBI - NP_001193658.1• AKT3 Chemiluminescent WBLot Kit Protocol

Application Details

Tested Applications:	IF, WB
Application Note:	Use Rockland Immunochemicals' Anti-AKT3 Chemiluminescent Kit for Western Blotting for detection of human AKT3 native and recombinant proteins by western blot. Expect a band approximately 56 kDa in size corresponding to AKT3 protein. This kit is useful for both western blotting and dot blotting methods. Please read the entire product insert prior to use.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
WB:	User Optimized

Formulation

Physical State:	n/a
Preservative:	Wash buffers MUST NOT contain SODIUM AZIDE or other inhibitors of peroxidase activity!
Stabilizer:	None


Shipping & Handling

Shipping Condition:	Dry Ice
----------------------------	---------

Storage Condition: See kit insert for complete instructions.


Expiration: See kit insert for complete instructions.

Images




Immunohistochemistry

Immunohistochemistry of Mouse Monoclonal anti AKT3 Antibody in Mouse Embryonic Kidney. Tissue: Mouse Liver. Fixation: FFPE buffered formalin 10% conc. Ag Retrieval: Heat, Citrate pH 6.2. Pressure Cooker. Primary antibody: anti-AKT3 at 2ug/ml for 1.5 hour @ room Temp. Secondary Ab: MOUSE ON MOUSE HRP POLYMER 45" RT.



Immunohistochemistry

Immunohistochemistry of Mouse Anti-AKT3 antibody. Tissue: human prostate carcinoma. A) AKT-3 antibody produced using CELLLine, B) AKT-3 antibody produced using roller bottle. Fixation: formalin fixed paraffin embedded. Antigen retrieval: not required. Primary antibody: AKT-3 antibody at 10 µg/mL for 1 h at RT. Secondary antibody: Peroxidase mouse secondary antibody at 1:10,000 for 1 h at RT. Localization: AKT3 is nuclear and occasionally cytoplasmic. Staining: AKT3 as precipitated brown signal with hematoxylin purple nuclear counterstain.




Western Blot

Western Blot of AKT isoform specific antibodies: 50 ng of recombinant GST-AKT1, -AKT2 and -AKT3 proteins were separated by SDS PAGE (4-20% gel) and transferred onto 0.2 µm nitrocellulose. Lane 1: GST Tagged recombinant AKT1. Lane 2: GST Tagged recombinant AKT2. Lane 3: GST Tagged recombinant AKT3. The membrane was blocked with Blocking Buffer (p/n MB-070) for 1 h at room temperature. Mouse anti-AKT1 (p/n 200-301-E75) at a dilution of 1 µg/mL in Blocking Buffer and incubated for 16 h at 4°C. Rabbit anti-Mouse IgG HRP (p/n 610-403-C46) secondary antibodies were applied at 0.05 µg/mL in Blocking Buffer and incubated for 1 h at room temperature followed by detection with FemtoMax™ chemiluminescent reagent (p/n FEMTOMAX-110). Predicted/Observed size: The expected size for all recombinant GST-AKT isoforms is 85 kDa. Arrowheads indicate the position of AKT isoform.


Western Blot

Western Blot of Mouse Anti-AKT3 antibody. Lane 1: C2C12. Lane 2: MEF#1. Lane 3: MEF#2. Lane 4: A549. Lane 5: Calu-1. Lane 6: PC3. Lane 7: HepG2. Lane 8: Jurkat. Lane 9: SKOV3. Lane 10: 293T. Load: 35 µg per lane. Primary antibody: AKT-3 antibody at 1:1000 for overnight at 4°C. Secondary antibody: Anti mouse secondary antibody at 1:20,000 for 1 h at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: 56 kDa for AKT3.



Western Blot

Western Blot of Mouse anti-AKT3 antibody. Lane 1: Control. Lane 2: Rapa. Lane 3: T50. Lane 4: T250. Lane 5: Control. Lane 6: Rapa. Lane 7: T50. Lane 8: T250. Lane 9: AKT3 null. Load: 35 µg per lane. Primary antibody: AKT-3 antibody at 1:1000 for overnight at 4°C. Secondary antibody: Anti mouse secondary antibody at 1:20,000 for 1 h at RT. Block: 5% BLOTTO overnight at 4°C. Predicted/Observed size: 56 kDa for AKT3.



Disclaimer

This material is subject to proprietary rights and is sold under license from ROCKLAND IMMUNOCHEMICALS Inc. This product is licensed for sale only for 'research-use' only. There is no implied license hereunder for any commercial use. FemtoMax[®] is a trademark of ROCKLAND IMMUNOCHEMICALS Inc. This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.