

Datasheet for KOA0350**Mouse CD40L ELISA Kit****Overview**

Description:	Mouse CD40L AccuSignal ELISA Kit - KOA0350
Item No.:	KOA0350
Size:	1 Kit
Applications:	ELISA
Reactivity:	Mouse
Range:	62.5 pg/ml - 4000 pg/ml

Product Details

Background:	CD40 ligand(CD40L) is a type II membrane protein of 261 amino acids on activated T cells that induces B cell proliferation and immunoglobulin secretion. It has homology with tumour necrosis factor-alpha and -beta, and has important functions in B-cell activation and differentiation. Human CD40L with 5 exons, is mapped to the proximal region of the mouse X chromosome on Xq26.3-27.1, and can be detected on T cells but is absent from B cells and monocytes. Since CD40L is expressed on platelets and released from them on activation, its predictive value as a marker for clinical outcome and the therapeutic effect of inhibition of glycoprotein IIb/IIIa receptor in patients with acute coronary syndromes was investigated. The soluble CD40L may be involved in the process of restenosis and that it exerts its effect by triggering a complex group of inflammatory reactions on endothelial and mononuclear cells. CD40L plays a central role in the pathophysiology of acute coronary syndromes, and has a role in the pathogenesis o...
Synonyms:	CD154, CD40 antigen ligand, CD40 ligand, CD40-L, CD40L_HUMAN, CD40LG, gp39, HIGM1, IGM, IMD3, soluble form, T B cell activating molecule, T BAM, T cell antigen Gp39, T-cell antigen Gp39, TNF related activation protein, TNF-related activation protein, TNFSF5, TrAP, Tumor necrosis factor (ligand) superfamily member 5, Tumor necrosis factor (ligand) superfamily, member 5 (hyper-IgM syndrome), Tumor necrosis factor ligand superfamily member 5
Detection Kit Type:	ELISA Kit
Detection Range:	62.5 pg/ml - 4000 pg/ml
Sensitivity:	<5 pg/ml

Target Details

Gene Name:	CD40LG
Reactivity:	Mouse
Immunogen:	Expression system for standard: NSO; Immunogen sequence: E61-L260
Purity/Specificity:	Natural and recombinant mouse CD40L. There is no detectable cross-reactivity with other relevant proteins.
Relevant Links:	• UniProtKB - P27548 • NCBI - NP_035746.2 • GeneID - 21947 • KOA0350 Protocol

Application Details

Tested Applications:	ELISA
Application Note:	Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1ml per well of the 4000pg/ml, 2000pg/ml, 1000pg/ml, 500pg/ml, 250pg/ml, 125pg/ml, 62.5pg/ml mouse CD40L standard solutions into the precoated 96-well plate. Add 0.1ml of the sample diluent buffer into the control well (Zero well). Add 0.1ml of each properly diluted sample of mouse cell culture supernates, serum or plasma (heparin, EDTA) to each empty well. It is recommended that each mouse CD40L standard solution and each sample be measured in duplicate.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	62.5pg/ml-4000pg/ml
Other:	DOE: @4°C 05/25/2026 @ -20°C 11/25/2026

Formulation

Concentration:	1 Kit
Anticoagulant:	Heparin Sodium

Shipping & Handling


Shipping Condition:	Wet Ice
---------------------	---------

Storage Condition:	Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.
Expiration:	See kit insert for complete instructions.

Images

ELISA

Mouse CD40L AccuSignal ELISA Kit standard curve. Assay Range: 62.5pg/ml-4000pg/ml. Sensitivity: <10pg/ml.



Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.