

## Datasheet for KOA0372

**Human ADAM8 ELISA Kit****Overview**

<b>Description:</b>	Human ADAM8 AccuSignal ELISA Kit - KOA0372
<b>Item No.:</b>	KOA0372
<b>Size:</b>	1 Kit
<b>Applications:</b>	ELISA
<b>Reactivity:</b>	Human
<b>Range:</b>	62.5 pg/ml - 4000 pg/ml

**Product Details**

<b>Background:</b>	A Disintegrin and metalloproteinase domain-containing protein 8 is an enzyme that in humans is encoded by the ADAM8 gene. ADAM8 is localized to chromosome 10q26.3. This gene encodes a member of the ADAM (a disintegrin and metalloproteinase domain) family. Members of the ADAM family, such as ADAM8, are cell surface proteases involved in remodeling of extracellular matrix, cell migration, and processing of membrane-bound signaling molecules. They are characterized by disintegrin and metalloprotease domains that confer adhesive properties and proteolytic activities, respectively. And the protein encoded by this gene may be involved in cell adhesion during neurodegeneration.
<b>Synonyms:</b>	A Disintegrin And Metalloproteinase domain 8, ADAM 8, ADAM 8 precursor, ADAM metallopeptidase domain 8, Adam8, ADAM8_HUMAN, CD 156, CD156, CD156a, CD156a antigen, CD156a antigen, Cell surface antigen MS2, Cell surface antigen MS2, Disintegrin and metalloproteinase domain-containing protein 8, Human leukocyte differentiation antigen, Human leukocyte differentiation antigen, Macrophage cysteine rich glycoprotein, MGC134985, MS 2, MS2
<b>Detection Kit Type:</b>	ELISA Kit
<b>Detection Range:</b>	62.5 pg/ml - 4000 pg/ml
<b>Sensitivity:</b>	<10 pg/ml

**Target Details**

<b>Gene Name:</b>	ADAM8
<b>Reactivity:</b>	Human

<b>Immunogen:</b>	Expression system for standard: NSO; Immunogen sequence: M1-P497
<b>Purity/Specificity:</b>	Natural and recombinant human ADAM8. There is no detectable cross-reactivity with other relevant proteins.
<b>Relevant Links:</b>	<ul style="list-style-type: none"><li>• <a href="#">UniProtKB - P78325</a></li><li>• <a href="#">NCBI - NP_001100.3</a></li><li>• <a href="#">GenID - 101</a></li><li>• <a href="#">KOA0372 Protocol</a></li></ul>

## Application Details


<b>Tested Applications:</b>	ELISA
<b>Application Note:</b>	Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1ml per well of the 4000pg/ml, 2000pg/ml, 1000pg/ml, 500pg/ml, 250pg/ml, 125pg/ml, 62.5pg/ml human ADAM8 standard solutions into the precoated 96-well plate. Add 0.1ml of the sample diluent buffer into the control well (Zero well). Add 0.1ml of each properly diluted sample of human cell culture supernates or serum to each empty well. It is recommended that each human ADAM8 standard solution and each sample be measured in duplicate.
<b>Assay Dilutions:</b>	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

## Formulation

## Shipping & Handling

<b>Shipping Condition:</b>	Wet Ice
<b>Storage Condition:</b>	Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.
<b>Expiration:</b>	See kit insert for complete instructions.

## Images

**ELISA**

Human ADAM8 AccuSignal ELISA Kit standard curve. Assay Range: 62.5pg/ml-4000pg/ml. Sensitivity: <10pg/ml.

**Disclaimer**

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.