

## Datasheet for KOA0384

**Human Cadherin-2 - N-Cadherin ELISA Kit****Overview**

<b>Description:</b>	Human Cadherin-2/N-Cadherin AccuSignal ELISA Kit - KOA0384
<b>Item No.:</b>	KOA0384
<b>Size:</b>	1 Kit
<b>Applications:</b>	ELISA
<b>Reactivity:</b>	Human
<b>Range:</b>	0.78 ng/ml - 50 ng/ml

**Product Details**

**Background:** Cadherin-2(CDH2), also known as neural cadherin(NCAD), is a protein that in humans is encoded by the CDH2 gene. It is a classical cadherin from the cadherin superfamily. This gene is mapped to 18q12.1. Cadherin-2 is expressed in the brain, skeletal and cardiac muscle. Cadherin-2 is commonly found in cancer cells and provides a mechanism for transendothelial migration. It is a calcium dependent cell-cell adhesion glycoprotein comprising five extracellular cadherin repeats, a transmembrane region and a highly conserved cytoplasmic tail. The protein functions during gastrulation and is required for establishment of left-right asymmetry. At certain central nervous system synapses, presynaptic to postsynaptic adhesion is mediated at least in part by this gene product.

<b>Synonyms:</b>	CDw325, CDHN, Neural cadherin, N-cadherin, CD325
<b>Detection Kit Type:</b>	ELISA Kit
<b>Detection Range:</b>	0.78 ng/ml - 50 ng/ml
<b>Sensitivity:</b>	<20 pg/ml

**Target Details**

<b>Gene Name:</b>	CDH2
<b>Reactivity:</b>	Human
<b>Immunogen:</b>	Expression system for standard: NSO; Immunogen sequence: D160-A724
<b>Purity/Specificity:</b>	Natural and recombinant human Cadherin-2. There is no detectable cross-reactivity with other relevant proteins.

<b>Relevant Links:</b>	<ul style="list-style-type: none"><li>• <a href="#">UniProtKB - P19022</a></li><li>• <a href="#">NCBI - NP_001295105.1</a></li><li>• <a href="#">GeneID - 1000</a></li><li>• <a href="#">KOA0384 Protocol</a></li></ul>
------------------------	---

## Application Details

<b>Tested Applications:</b>	ELISA
<b>Application Note:</b>	Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1ml per well of the 50ng/ml, 25ng/ml, 12.5ng/ml, 6.25ng/ml, 3.12ng/ml, 1.56ng/ml, 0.78ng/ml human Cadherin-2 standard solutions into the precoated 96-well plate. Add 0.1ml of the sample diluent buffer into the control well (Zero well). Add 0.1ml of each properly diluted sample of human cell culture supernates, serum or plasma(heparin, EDTA) to each empty well. We recommend that each human Cadherin-2 standard solution and each sample is measured in duplicate.
<b>Assay Dilutions:</b>	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

## Formulation

<b>Anticoagulant:</b>	Heparin Sodium
-----------------------	----------------


## Shipping & Handling

<b>Shipping Condition:</b>	Wet Ice
<b>Storage Condition:</b>	Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.
<b>Expiration:</b>	See kit insert for complete instructions.

## Images

**ELISA**

Human Cadherin-2/N-Cadherin AccuSignal ELISA Kit  
standard curve. Assay Range: 0.78ng/ml-50ng/ml.  
Sensitivity: <20pg/ml.

**Disclaimer**

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.