

## Datasheet for KOA0453

**Human GDF15 ELISA Kit****Overview**

<b>Description:</b>	Human GDF-15 AccuSignal ELISA Kit - KOA0453
<b>Item No.:</b>	KOA0453
<b>Size:</b>	1 Kit
<b>Applications:</b>	ELISA
<b>Reactivity:</b>	Human
<b>Range:</b>	31.2 pg/ml - 2000 pg/ml

**Product Details**

<b>Background:</b>	GDF-15(Growth differentiation factor 15),also known as TGF-PL, MIC-1, PDF, PLAB, and PTGFB, is a protein belonging to the transforming growth factor beta superfamily that has a role in regulating inflammatory and apoptotic pathways in injured tissues and during disease processes. Using FISH, the MIC1 gene is mapped to 19p13.2-p13.1. Its expression in liver can be significantly up-regulated in during injury of organs such as liver, kidney, heart and lung. GDF15 showed increased expression and secretion during erythroblast maturation. GDF15 functions as an anti-inflammatory cytokine by directly interfering with chemokine signaling and integrin activation.
<b>Synonyms:</b>	Growth/differentiation factor 15, Macrophage inhibitory cytokine 1, MIC 1, NAG-1, NRG-1, NSAID-activated gene 1 protein, NSAID-regulated gene 1 protein, PDF, PLAB, Placental bone morphogenetic protein, Placental TGF-beta, Prostate differentiation factor, PROSTATE-DERIVED FACTOR, PTGFB
<b>Detection Kit Type:</b>	ELISA Kit
<b>Detection Range:</b>	31.2 pg/ml - 2000 pg/ml
<b>Sensitivity:</b>	<10 pg/ml

**Target Details**

<b>Gene Name:</b>	GDF15
<b>Reactivity:</b>	Human
<b>Immunogen:</b>	Expression system for standard: CHO; Immunogen sequence: A197-I308

**Purity/Specificity:** Natural and recombinant human GDF-15. There is no detectable cross-reactivity with other relevant proteins.

**Relevant Links:**

- [UniProtKB - Q99988](#)
- [NCBI - NP\\_004855.2](#)
- [GeneID - 9518](#)
- [KOA0453 Protocol](#)

## Application Details

**Tested Applications:** ELISA

**Application Note:** Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1ml per well of the 2000pg/ml, 1000pg/ml, 500pg/ml, 250pg/ml, 125pg/ml, 62.5pg/ml, 31.2pg/ml human GDF-15 standard solutions into the pre-coated 96-well plate. Add 0.1ml of the sample diluent buffer into the control well (Zero well). Add 0.1ml of each properly diluted sample of human cell culture supernates, serum or plasma (heparin, EDTA) to each empty well. It is recommended that each human GDF-15 standard solution and each sample be measured in duplicate.

**Assay Dilutions:** All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

**ELISA:** 31.2pg/ml-2000pg/ml

**Other:**  
DOE:  
2-8° C - 01/29/2025  
-20° C - 07/29/2025

## Formulation

**Concentration:** 1 Kit

**Anticoagulant:** Heparin Sodium

## Shipping & Handling


**Shipping Condition:** Wet Ice

**Storage Condition:** Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.

**Expiration:** See kit insert for complete instructions.

## Images

### Human GDF-15 ELISA Kit



### ELISA

Human GDF-15 AccuSignal ELISA Kit standard curve. Assay Range: 31.2pg/ml-2000pg/ml. Sensitivity: <10pg/ml.

## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.