

Datasheet for KOA0511**Human MPO ELISA Kit****Overview**

Description:	Human MPO AccuSignal ELISA Kit - KOA0511
Item No.:	KOA0511
Size:	1 Kit
Applications:	ELISA
Reactivity:	Human
Range:	312 pg/ml - 20000 pg/ml

Product Details

Background:	Myeloperoxidase(MPO) is a mammalian phagocyte hemoprotein thought to primarily mediate host defense reactions. It is abundantly expressed in neutrophils and secreted during their activation. Myeloperoxidase is part of the host defense system of human polymorphonuclear leukocytes, responsible for microbicidal activity against a wide range of organisms. It is located in the nucleus as well as in the cytoplasm. Intranuclear MPO may help to protect DNA against damage resulting from oxygen radicals produced during myeloid cell maturation and function. The standard product used in this kit is the product of gene recombination, consisting of 697 (A49-S745) amino acids with the molecular mass of 80KDa.
Synonyms:	84 kDa myeloperoxidase, 89 kDa myeloperoxidase, MPO, MPO Myeloperoxidase, Myeloperoxidase, Myeloperoxidase heavy chain, Myeloperoxidase light chain, PERM_HUMAN
Detection Kit Type:	ELISA Kit
Detection Range:	312 pg/ml - 20000 pg/ml
Sensitivity:	<10 pg/ml

Target Details

Gene Name:	MPO
Reactivity:	Human
Immunogen:	Expression system for standard: NSO; Immunogen sequence: A49-S745
Purity/Specificity:	Natural and recombinant human MPO. There is no detectable cross-reactivity with other relevant proteins.

Relevant Links:	<ul style="list-style-type: none">• UniProtKB - P05164• NCBI - NP_000241.1• GeneID - 4353• KOA0511 Protocol
------------------------	--

Application Details

Tested Applications:	ELISA
Application Note:	Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1ml per well of the 20000pg/ml, 10000pg/ml, 5000pg/ml, 2500pg/ml, 1250pg/ml, 625pg/ml, 312pg/ml human MPO standard solutions into the precoated 96-well plate. Add 0.1ml of the sample diluent buffer into the control well (Zero well). Add 0.1ml of each properly diluted sample of human cell culture supernates, cell lysate, tissue lysate, serum, plasma(heparin, EDTA), saliva or urine to each empty well. It is recommended that each human MPO standard solution and each sample be measured in duplicate.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
Other:	DOE: @4°C 10/16/2026 @-20°C 04/16/2027


Formulation

Concentration:	1 Kit
Anticoagulant:	Heparin Sodium

Shipping & Handling

Shipping Condition:	Wet Ice
Storage Condition:	Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.
Expiration:	See kit insert for complete instructions.

Images



ELISA

Human MPO AccuSignal ELISA Kit standard curve. Assay Range: 312pg/ml-20000pg/ml. Sensitivity: <10pg/ml.

References

- Qu X et al. Indole-3-acetic acid ameliorates dextran sulfate sodium-induced colitis via the ERK signaling pathway. *Arch Pharm Res.* (2024)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.