

Datasheet for KOA0513

Mouse Neprilysin - CD10 ELISA Kit**Overview**

Description:	Mouse Neprilysin/CD10 AccuSignal ELISA Kit - KOA0513
Item No.:	KOA0513
Size:	1 Kit
Applications:	ELISA
Reactivity:	Mouse
Range:	93.7 pg/ml - 6000 pg/ml

Product Details

Background:	Neprilysin, also known as membrane metallo-endopeptidase, neutral endopeptidase (NEP), CD10, and common acute lymphoblastic leukemia antigen (CALLA), is a zinc-dependent metalloprotease enzyme that degrades a number of small secreted peptides, most notably the amyloid beta peptide whose abnormal misfolding and aggregation in neural tissue has been implicated as a cause of Alzheimer's disease, this gene is localized to human chromosome 3 by study of somatic cell hybrids and regionalized the location to 3q21-q27 by in situ hybridization. By cDNA transfection analysis, Neprilysin is confirmed as a functional neutral endopeptidase of the type that has previously been called enkephalinase. Neprilysin has also been called atriopeptidase. Atriopeptidase specifically degrades atrial natriuretic factor. A specific enzyme inhibitor was developed and reported that it had effects similar to those of low-dose ANF infusion. These effects include diuresis, natriuresis, vasodilatation, and suppression of the renin-angiotensin-aldosterone system.
Synonyms:	Atriopeptidase, CALLA, CD10, CD10 antigen, Common acute lymphocytic leukemia antigen, Enkephalinase, EPN, gp100, Membrane metallo endopeptidase, Membrane metallo endopeptidase variant 1, Membrane metallo endopeptidase variant 2, Membrane metalloendopeptidase, Membrane metalloendopeptidase neutral endopeptidase, Membrane metalloendopeptidase variant 1, Membrane metalloendopeptidase variant 2, NEP, NEP_HUMAN, Neprilysin, Neutral endopeptidase, SFE, Skin fibroblast elastase
Detection Kit Type:	ELISA Kit
Detection Range:	93.7 pg/ml - 6000 pg/ml
Sensitivity:	<10 pg/ml

Target Details

Gene Name:	MME
Reactivity:	Mouse
Immunogen:	Expression system for standard: NSO; Immunogen sequence: Y52-W750
Purity/Specificity:	Natural and recombinant mouse CD10. There is no detectable cross-reactivity with other relevant proteins.
Relevant Links:	• UniProtKB - Q61391 • NCBI - NP_001276391.1 • GeneID - 17380 • KOA0513 Protocol

Application Details

Tested Applications:	ELISA
Application Note:	Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1ml per well of the 6000pg/ml, 3000pg/ml, 1500pg/ml, 750pg/ml, 419pg/ml, 187.5pg/ml, 93.7pg/ml mouse CD10 standard solutions into the precoated 96-well plate. Add 0.1ml of the sample diluent buffer into the control well (Zero well). Add 0.1ml of each properly diluted sample of mouse cell culture supernates, serum or plasma(heparin, EDTA) to each empty well. It is recommended that each mouse CD10 standard solution and each sample be measured in duplicate.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

Formulation


Anticoagulant:	Heparin Sodium
----------------	----------------

Shipping & Handling

Shipping Condition:	Wet Ice
Storage Condition:	Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.

Expiration: See kit insert for complete instructions.

Images



ELISA

Mouse Neprilysin/CD10 AccuSignal ELISA Kit standard curve.
Assay Range: 93.7pg/ml-6000pg/ml. Sensitivity: <10pg/ml.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.