

Datasheet for KOA0518**Human TNFSF4 - OX40L ELISA Kit****Overview**

Description:	Human TNFSF4/OX40L AccuSignal ELISA Kit - KOA0518
Item No.:	KOA0518
Size:	1 Kit
Applications:	ELISA
Reactivity:	Human
Range:	156 pg/ml - 10000 pg/ml

Product Details

Background:	OX40L is the ligand for CD134 and is expressed on such cells as DC2s enabling amplification of Th2 cell differentiation. OX40L has also been designated CD252. By analysis of an interspecific backcross, the mouse <i>Tnfsf4</i> gene is mapped to chromosome 1. Using fluorescence in situ hybridization, they localized the human TNFSF4 gene to 1q25, a region sharing homology of synteny with the portion of mouse chromosome 1 containing the <i>Tnfsf4</i> gene. OX40L strongly inhibited the generation of IL-10-producing Tr1 cells induced by two physiologic stimuli, the inducible costimulatory ligand and immature dendritic cells. In addition, OX40L strongly inhibited IL-10 production and suppressive function of differentiated IL-10-producing Tr1 cells. <i>Tnfsf4</i> underlies Ath1 in mice and that polymorphisms in its human homolog TNFSF4 increase the risk of myocardial infarction in humans.
Synonyms:	CD 134L, CD 252, CD134, CD134 ligand, CD134L, CD252, CD252 antigen, glycoprotein 34 kd, Glycoprotein Gp34, GP34, OX-40L, OX40 antigen ligand, OX40 ligand, OX40L, TAX transcriptionally-activated glycoprotein 1, TNFL4_HUMAN, <i>Tnfsf4</i> , Tumor necrosis factor (ligand) superfamily member 4, Tumor necrosis factor ligand superfamily member 4, TXGP1
Detection Kit Type:	ELISA Kit
Detection Range:	156 pg/ml - 10000 pg/ml
Sensitivity:	<10 pg/ml

Target Details

Gene Name:	TNFSF4
Reactivity:	Human

Immunogen:	Expression system for standard: NSO; Immunogen sequence: Q51-L183
Purity/Specificity:	Natural and recombinant human TNFSF4. There is no detectable cross-reactivity with other relevant proteins.
Relevant Links:	• UniProtKB - P23510 • NCBI - NP_001284491.1 • GenelD - 7292 • KOA0518 Protocol

Application Details

Tested Applications:	ELISA
Application Note:	Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1ml per well of the 10000pg/ml, 5000pg/ml, 2500pg/ml, 1250pg/ml, 625pg/ml, 313pg/ml, 156pg/ml human TNFSF4 standard solutions into the precoated 96-well plate. Add 0.1ml of the sample diluent buffer into the control well (Zero well). Add 0.1ml of each properly diluted sample of human cell culture supernates, serum or plasma (heparin, EDTA) to each empty well. It is recommended that each human TNFSF4 standard solution and each sample be measured in duplicate.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	156pg/ml-10000pg/ml
Other:	DOE: 2-8° C - 06/30/19 -20° C - 12/29/19

Formulation


Concentration:	1 Kit
Anticoagulant:	Heparin Sodium

Shipping & Handling

Shipping Condition:	Wet Ice
Storage Condition:	Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.

Expiration: See kit insert for complete instructions.

Images



ELISA

Human TNFSF4/OX40L AccuSignal ELISA Kit standard curve. Assay Range: 156pg/ml-10000pg/ml. Sensitivity: <10pg/ml.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.