

Datasheet for KOA0629

Mouse TGF beta 3 ELISA Kit**Overview**

Description:	Mouse TGF-beta 3 AccuSignal ELISA Kit - KOA0629
Item No.:	KOA0629
Size:	1 Kit
Applications:	ELISA
Reactivity:	Mouse
Range:	31.2 pg/ml - 2000 pg/ml

Product Details

Background:	Transforming growth factor beta 3(TGF-beta 3) is a type of protein, known as a cytokine, which is involved in cell differentiation, embryogenesis and development. It belongs to a large family of cytokines called the Transforming growth factor beta superfamily. TGF-beta 3 is believed to regulate molecules involved in cellular adhesion and extracellular matrix(ECM) formation during the process of palate development. Without TGF-beta 3, mammals develop a deformity known as a cleft palate. This is caused by failure of epithelial cells in both sides of the developing palate to fuse. TGF-beta 3 also plays an essential role in controlling the development of lungs in mammals, by also regulating cell adhesion and ECM formation in this tissue, and controls wound healing by regulating the movements of epidermal and dermal cells in injured skin. TGF-beta 3 activates Lef1 in the absence of beta-catenin(CTNNB1) via nuclear phospho-Smad2 and Smad4.
Synonyms:	ARVD, ARVD1, FLJ16571, MGC105479, MGC118722, prepro-transforming growth factor beta-3, RNHF, TGF beta 3, TGF beta3, TGF-beta-3, TGFB 3, TGFB3, TGFB3_HUMAN, transforming growth factor beta 3, Transforming growth factor beta-3
Detection Kit Type:	ELISA Kit
Detection Range:	31.2 pg/ml - 2000 pg/ml
Sensitivity:	<10 pg/ml

Target Details

Gene Name:	TGFB3
Reactivity:	Mouse

Immunogen:	Expression system for standard: sf21; Immunogen sequence: A301-S412
Purity/Specificity:	Natural and recombinant mouse TGF-beta 3. There is no detectable cross-reactivity with other relevant proteins.
Relevant Links:	• UniProtKB - P17125 • NCBI - NP_033394.2 • GenelD - 21809 • KOA0629 Protocol

Application Details

Tested Applications:	ELISA
Application Note:	Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1ml per well of the 2000pg/ml, 1000pg/ml, 500pg/ml, 250pg/ml, 125pg/ml, 62.5pg/ml, 31.2pg/ml mouse TGF-beta 3 standard solutions into the precoated 96-well plate. Add 0.1ml of the sample diluent buffer into the control well (Zero well). Add 0.1ml of each properly diluted sample of activated mouse cell culture supernates, serum, plasma(EDTA) or urine to each empty well. It is recommended that each mouse TGF-beta 3 standard solution and each sample be measured in duplicate.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	31.2pg/ml-2000pg/ml
Other:	DOE: 2-8° C - 02/16/2024 -20° C - 08/16/2024

Formulation


Concentration:	1 Kit
Anticoagulant:	Sodium EDTA

Shipping & Handling

Shipping Condition:	Wet Ice
Storage Condition:	Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.

Expiration: See kit insert for complete instructions.

Images



ELISA

Mouse TGF-beta 3 AccuSignal ELISA Kit standard curve.
Assay Range: 31.2pg/ml-2000pg/ml. Sensitivity: <10pg/ml.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.