

Datasheet for KOA0686**Mouse Periostin - OSF2 ELISA Kit****Overview**

Description:	Mouse Periostin/OSF2 AccuSignal ELISA Kit - KOA0686
Item No.:	KOA0686
Size:	1 Kit
Applications:	ELISA
Reactivity:	Mouse
Range:	93.7 pg/ml - 6000 pg/ml

Product Details

Background:	Periostin, also known as OSF2, is a protein that in humans is encoded by the POSTN gene. The International Radiation Hybrid Mapping Consortium mapped the POSTN gene to chromosome 13. Periostin functions as a ligand for alpha-V/beta-3 and alpha-V/ beta-5 integrins to support adhesion and migration of epithelial cells. It is found that periostin was overexpressed by the majority of human primary breast cancers examined. After myocardial infarction, periostin-induced cardiomyocyte cell cycle reentry and mitosis were associated with improved ventricular remodeling and myocardial function, reduced fibrosis and infarct size, and increase angiogenesis.
Synonyms:	OSF-2, Osteoblast-specific factor 2, Periostin, PN, POSTN, POSTN_HUMAN
Detection Kit Type:	ELISA Kit
Detection Range:	93.7 pg/ml - 6000 pg/ml
Sensitivity:	<10 pg/ml

Target Details

Gene Name:	POSTN
Reactivity:	Mouse
Immunogen:	Expression system for standard: NSO; Immunogen sequence: N24-Q836
Purity/Specificity:	Natural and recombinant mouse Periostin. There is no detectable cross-reactivity with other relevant proteins.

Relevant Links:

- [UniProtKB - Q62009](#)
- [NCBI - NP_001185694.1](#)
- [GeneID - 50706](#)
- [KOA0686 Protocol](#)

Application Details

Tested Applications: ELISA

Application Note: Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1ml per well of the 6000pg/ml, 3000pg/ml, 1500pg/ml, 750pg/ml, 419pg/ml, 187.5pg/ml, 93.7pg/ml mouse Periostin standard solutions into the precoated 96-well plate. Add 0.1ml of the sample diluent buffer into the control well (Zero well). Add 0.1ml of each properly diluted sample of mouse cell culture supernates, serum or plasma (heparin, EDTA) to each empty well. It is recommended that each mouse Periostin standard solution and each sample be measured in duplicate.

Assay Dilutions: All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

Formulation

Anticoagulant: Heparin Sodium

Shipping & Handling


Shipping Condition: Wet Ice

Storage Condition: Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.

Expiration: See kit insert for complete instructions.

Images

Mouse OSF2 ELISA Kit



ELISA

Mouse Periostin/OSF2 ELISA Kit standard curve.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.