

Datasheet for KOA0709

Human CD13 - Aminopeptidase N ELISA Kit**Overview**

Description:	Human CD13/Aminopeptidase N AccuSignal ELISA Kit - KOA0709
Item No.:	KOA0709
Size:	1 Kit
Applications:	ELISA
Reactivity:	Human
Range:	312 pg/ml - 20,000 pg/ml

Product Details

Background:	Alanine aminopeptidase, also known as ANPEP or CD13, is an enzyme that is used as a biomarker to detect damage to the kidneys, and that may be used to help diagnose certain kidney disorders. It is mapped to 15q26.1. Aminopeptidase N is located in the small-intestinal and renal microvillar membrane, and also in other plasma membranes. In the small intestine, Aminopeptidase N plays a role in the final digestion of peptides generated from hydrolysis of proteins by gastric and pancreatic proteases, and it is also thought to be involved in the metabolism of regulatory peptides by diverse cell types, including small intestinal and renal tubular epithelial cells, macrophages, granulocytes, and synaptic membranes from the CNS. Petrovic et al showed that CD13 was required for endothelial cell invasion in response to bradykinin. Inhibition of CD13 abrogated internalization of bradykinin receptor B2 and reduced endothelial cell motility.
Synonyms:	Alanyl (membrane) aminopeptidase, Alanyl aminopeptidase, Alanyl membrane aminopeptidase, Aminopeptidase M, Aminopeptidase N, AMPN_HUMAN, ANPEP, AP M, AP N, AP-M, AP-N, APN, CD 13, CD13, CD13 antigen, gp150, hAPN, Lap 1, Lap1, Microsomal aminopeptidase, Myeloid plasma membrane glycoprotein CD13, p150, PEPN
Detection Kit Type:	ELISA Kit
Detection Range:	312 pg/ml - 20,000 pg/ml
Sensitivity:	<10 pg/ml

Target Details

Gene Name:	ANPEP
-------------------	-------

Reactivity:	Human
Immunogen:	Expression system for standard: NSO; Immunogen sequence: K69-K967
Purity/Specificity:	Natural and recombinant human CD13. There is no detectable cross-reactivity with other relevant proteins.
Relevant Links:	<ul style="list-style-type: none">• UniProtKB - P15144• NCBI - NP_001141.2• GenelD - 290• KOA0709 Protocol

Application Details

Tested Applications:	ELISA
Application Note:	Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1ml per well of the 20,000pg/ml, 10,000pg/ml, 5000pg/ml, 2500pg/ml, 1250pg/ml, 625pg/ml, 312pg/ml human CD13 standard solutions into the precoated 96-well plate. Add 0.1ml of the sample diluent buffer into the control well (Zero well). Add 0.1ml of each properly diluted sample of human cell culture supernates, serum or plasma(heparin, EDTA) to each empty well. It is recommended that each human CD13 standard solution and each sample be measured in duplicate.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

Formulation


Anticoagulant:	Heparin Sodium
-----------------------	----------------

Shipping & Handling

Shipping Condition:	Wet Ice
Storage Condition:	Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.
Expiration:	See kit insert for complete instructions.

Images

Human CD13 ELISA Kit



ELISA

Human CD13/Aminopeptidase N ELISA Kit standard curve.

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.