

## Datasheet for KOA0875

**Mouse PD-L1 - B7-H1 ELISA Kit****Overview**

<b>Description:</b>	Mouse PD-L1/B7-H1 AccuSignal ELISA Kit - KOA0875
<b>Item No.:</b>	KOA0875
<b>Size:</b>	1 Kit
<b>Applications:</b>	ELISA
<b>Reactivity:</b>	Mouse
<b>Range:</b>	31.2 pg/ml - 2000 pg/ml

**Product Details**

<b>Background:</b>	Programmed death-ligand 1 (PD-L1), also known as cluster of differentiation 274 (CD274) or B7 homolog 1 (B7-H1), is a protein that in humans is encoded by the CD274 gene. PD-L1 is a 40kDa type 1 transmembrane protein that has been speculated to play a major role in suppressing the immune system during particular events such as pregnancy, tissue allografts, autoimmune disease and other disease states such as hepatitis. Normally the immune system reacts to foreign antigens where there is some accumulation in the lymph nodes or spleen which triggers a proliferation of antigen-specific CD8+ T cell. The formation of PD-1 receptor / PD-L1 or B7.1 receptor /PD-L1 ligand complex transmits an inhibitory signal which reduces the proliferation of these CD8+ T cells at the lymph nodes and supplementary to that PD-1 is also able to control the accumulation of foreign antigen specific T cells in the lymph nodes through apoptosis which is further mediated by a lower regulation of the gene Bcl-2.
<b>Synonyms:</b>	B7 H, B7H, B7 H1, B7-H1, B7H1, CD 274, CD274, PD L1, PD-L1, PDL 1, PDL1
<b>Detection Kit Type:</b>	ELISA Kit
<b>Detection Range:</b>	31.2 pg/ml - 2000 pg/ml
<b>Sensitivity:</b>	<10 pg/ml

**Target Details**

<b>Gene Name:</b>	CD274
<b>Reactivity:</b>	Mouse
<b>Immunogen:</b>	Expression system for standard: NSO; Immunogen sequence: F19-H239

**Purity/Specificity:** Natural and recombinant mouse PD-L1. There is no detectable cross-reactivity with other relevant proteins.

**Relevant Links:**

- [UniProtKB - Q9EP73](#)
- [NCBI - NP\\_068693.1](#)
- [GeneID - 60533](#)
- [KOA0875 Protocol](#)

## Application Details

**Tested Applications:** ELISA

**Application Note:** Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1ml per well of the 2000pg/ml, 1000pg/ml, 500pg/ml, 250pg/ml, 125pg/ml, 62.5pg/ml, 31.2pg/ml mouse PD-L1 standard solutions into the pre-coated 96-well plate. Add 0.1ml of the sample diluent buffer into the control well (Zero well). Add 0.1ml of each properly diluted sample of mouse cell culture supernates, serum or plasma (heparin, EDTA) to each empty well. It is recommended that each mouse PD-L1 standard solution and each sample be measured in duplicate.

**Assay Dilutions:** All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

**ELISA:** 31.2 pg/ml-2000 pg/ml

**Other:**  
DOE:  
2-8° C - 08/17/2022  
-20° C - 02/15/2023

## Formulation

**Concentration:** 1 Kit

**Anticoagulant:** Heparin Sodium


## Shipping & Handling

**Shipping Condition:** Wet Ice

**Storage Condition:** Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.

**Expiration:** See kit insert for complete instructions.

## Images



### ELISA

Mouse PD-L1/B7-H1 ELISA Kit standard curve.

## References

- Wang Y et al. Immuno-protective vesicle-crosslinked hydrogel for allogenic transplantation. *Nat Commun.* (2024)

## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.