

Datasheet for PA00-03-1000**Protein A Peroxidase Conjugated****Overview**

Description:	Protein A Peroxidase Conjugated - PA00-03-1000
Item No.:	PA00-03-1000
Size:	1 mg
Applications:	Dot Blot, ELISA, WB
Origin:	Staphylococcus aureus

Product Details

Background:	Protein A is a cell wall protein originating from the bacterium <i>Staphylococcus aureus</i>. Protein A binds to immunoglobulins through interaction with the IgG heavy chain within the Fc region. Affinity to Fc may vary depending on IgG species origins or subclasses. Protein A is predicted 42kDa based on sedimentation data and observed to run ~55-56kDa on SDS-Page. This high affinity property makes Protein A essential in the large scale purification of antibodies. Horseradish Peroxidase (HRP) is an enzyme that utilize organic peroxide compounds as electron donors. Naturally provides protection for plants against pathogens, but can be utilized in molecular biology to convert various substrates to detectable compounds (such as in Western Blotting and ELISAs).
Synonyms:	HRP, peroxidase, Horseradish peroxidase protein A
Species of Origin:	<i>Staphylococcus aureus</i>
Conjugate:	Peroxidase (HRP)

Target Details

Purity/Specificity:	Protein A Peroxidase was prepared from chromatographically pure Protein A. Protein A Peroxidase was assayed by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase and anti-Protein A. No reaction was observed against anti-Protein G.
Relevant Links:	• SDS

Application Details

Tested Applications:	Dot Blot
Suggested Applications:	ELISA, WB (Based on references)
Application Note:	Protein A Peroxidase Conjugate has been tested by dot blot and is a useful reagent in Western Blotting and ELISA experiments. Protein A Peroxidase can be utilized as a pseudo-secondary detection reagent when used in conjunction with an IgG-based primary antibody and appropriate substrate (such as TMB-1000 or Femtomax-110).
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:20,000 - 1:200,000
IHC:	1:1,000 - 1:5,000
WB:	1:10,000 - 1:40,000

Formulation

Physical State:	Lyophilized
Concentration:	1.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!
Stabilizer:	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Reconstitution Volume:	1.0 mL
Reconstitution Buffer:	Restore with deionized water (or equivalent)

Shipping & Handling


Shipping Condition:	Ambient
Storage Condition:	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. Protein A Peroxidase conjugated is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Images

ELISA target and detector	NDO-LID		
	IgM	IgG	IgG/M (protein A)
Sensitivity	78.0 (73.0–83.0)	81.6 (77.0–86.1)	86.3 (82.2–90.3)
Specificity	97.0 (94.0–100)	95.6 (92.0–99.4)	93.6 (93.0–100)
PPV	98.3 (96.5–100)	97.6 (95.5–99.7)	98.1 (96.3–100)
NPV	66.7 (60.0–73.5)	70.2 (63.4–77.1)	76.1 (69.5–82.8)

ELISA

Diagnostic accuracy of detecting IgG and IgG/M antibodies against natural octyl disaccharide-leprosy IDRI diagnostic (NDO-LID) using HRP anti-human IgG (p/n 609-1316), HRP anti-human IgM (p/n 609-4331), and HRP Protein A (p/n PA 00-03). ELISA = enzyme-linked immunosorbent assay; NPV = negative predictive value; PPV = positive predictive value. Results are shown as percent, with confidence intervals in parentheses. Table 5. PMID: 29141725.



Dot Blot

Dot Blot of PEROXIDASE Conjugated Protein A. Lane 1: Human IgG. Lane 2: Golden Syrian Hamster IgG. Lane 3: Sheep IgG. Load: 3-fold serial dilution starting at 200 ng. Primary Antibody: none. Secondary antibody: PEROXIDASE Conjugated Protein A at 1:1000 for 60 min at RT. Block: 1% BSA-TTBS 30 min at RT.

References

- Zhang W et al. GCAF(TM251) regulates lysosome biogenesis by activating the mannose-6-phosphate pathway. *Nat Commun.* (2022)
- Munoz et al. Comparison of Enzyme-Linked Immunosorbent Assay Using Either Natural Octyl Disaccharide-Leprosy IDRI Diagnostic or Phenolic Glycolipid-I Antigens for the Detection of Leprosy Patients in Colombia. *The American Journal of Tropical Medicine and Hygiene* (2018)
- Serrano-Coll H et al. Anti-natural octyl disaccharide-leprosy IDRI diagnostic (NDO-LID) antibodies as indicators of leprosy reactions and neuritis. *Trans R Soc Trop Med Hyg.* (2017)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.