

Datasheet for PG00-03**Protein G Peroxidase Conjugated****Overview**

Description:	Protein G Peroxidase Conjugated - PG00-03
Item No.:	PG00-03
Size:	1 mg
Applications:	Dot Blot, ELISA
Origin:	Streptococcus sp.

Product Details

Background:	Protein G is a surface protein of two groups of Streptococcal bacteria that has the ability to bind immunoglobulins. Similar to Protein A, but with slightly different specificity, Protein G is an important agent in the purification of proteins due to its ability to bind the Fc region. While native Protein G binds to albumin, recombinant Protein G is designed to contain only immunoglobulin binding domains to ensure the maximum specific IgG binding capacity. Horseradish Peroxidase (HRP) is an enzyme that utilize organic peroxide compounds as electron donors. Naturally provides protection for plants against pathogens, but can be utilized in molecular biology to convert various substrates to detectable compounds (such as in Western Blotting and ELISAs).
Synonyms:	ProG, Streptococcus G protein, Protein G HRP, peroxidase, Horseradish peroxidase
Species of Origin:	Streptococcus sp.
Conjugate:	Peroxidase (HRP)
F/P Ratio:	1.54

Target Details

Purity/Specificity:	Protein G Peroxidase was prepared from chromatographically pure recombinant Protein G. Protein G Peroxidase was assayed by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase and anti-Protein G. No reaction was observed against anti-Protein A.
Relevant Links:	• UniProtKB - P19909 • PG00-03 SDS

Application Details

Tested Applications:	Dot Blot, ELISA
Application Note:	Protein G Peroxidase has been tested by ELISA and dot blot and is a useful reagent in Western Blotting and ELISA experiments. Protein G Peroxidase can be utilized as a pseudo-secondary detection reagent when used in conjunction with an IgG-based primary antibody and appropriate substrate (such as TMB-1000 or Fentomax-110).
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
ELISA:	1:20,000 - 1:200,000
IHC:	1:1,000 - 1:5,000
WB:	1:10,000 - 1:40,000


Formulation

Physical State:	Lyophilized
Concentration:	1.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!
Stabilizer:	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Reconstitution Volume:	1.0 mL
Reconstitution Buffer:	Restore with deionized water (or equivalent)

Shipping & Handling

Shipping Condition:	Ambient
Storage Condition:	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. Protein G Peroxidase conjugated is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.

Images



Dot Blot

Dot Blot of Protein G Peroxidase Conjugated. Antigen 1: Human IgG. Antigen 2: Rat IgG. Antigen 3: Dog IgG. Load: 200ng followed by a 3-fold serial dilution. Primary antibody: none. Secondary antibody: Protein G Peroxidase Conjugated at 1:1,000 for 60 min at RT. Block: MB-070 for 60 min at RT.

References

- Gotto S. et al. Studies on the mechanism of the immunosuppression caused by bovine mycoplasmosis and the development of its novel control method. *Hokkaido University Collection of Scholarly and Academic Papers*. (2020)
- Goto S, Konnai S, Hirano Y, et al. Upregulation of PD-L1 Expression by Prostaglandin E2 and the Enhancement of IFN- γ by Anti-PD-L1 Antibody Combined With a COX-2 Inhibitor in Mycoplasma bovis Infection. *Front Vet Sci*. (2020)
- Angeles, JMM et al. Field Evaluation of Recombinant Antigen ELISA in Detecting Zoonotic Schistosome Infection Among Water Buffaloes in Endemic Municipalities in the Philippines. *Frontiers in Veterinary Science* (2020)
- Angeles, JMM et al. Utilization of ELISA using thioredoxin peroxidase-1 and tandem repeat proteins for diagnosis of Schistosoma japonicum infection among water buffaloes. *PLoS Neglected Tropical Diseases* (2012)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.