

Datasheet for S000-06**Streptavidin Cy5 Conjugated****Overview**

Description:	Streptavidin Cy5 Conjugated - S000-06
Item No.:	S000-06
Size:	1 mg
Applications:	Dot Blot, WB, ELISA, EM, FISH, Functional Assay, IF, IHC, Microarray, Multiplex

Product Details

Background:	Streptavidin is isolated from bacteria, <i>Streptomyces avidinii</i> , and has an exceptionally high binding affinity for B7 (biotin). Rockland offers streptavidin in unconjugated and conjugated forms for common immunoassays including ELISA, western blotting, immunohistochemistry. Streptavidin is a tetrameric protein capable of binding 4 biotin groups to each molecule of streptavidin. While streptavidin has identical binding properties as avidin, it lacks the glycoprotein portion of the molecule and therefore shows less non-specific binding. Streptavidin is a slightly smaller molecule with a molecular weight of approximately 53.6 kDa. The sequence of avidin only shows 30% homology with streptavidin, and anti-avidin and anti-streptavidin antibodies are not immunologically cross reactive. Rockland conjugates a broad group of secondary antibodies to many of the classic fluorescent markers including fluorescein, rhodamine, Texas Red, CyDyes™ and Phycoerythrin (RPE). All of the conjugates are ideal for various immunofluorescence based assays including fluorescent western blotting, immunofluorescence microscopy, FLISA, and more. Rockland also produces many next generation fluorochrome dyes designed for detection of primary antibodies in multiplex, multi-color analysis.
Synonyms:	SA, S avidin, streptococcus avidin, Streptavidin Cy5 Conjugated, Cy5 conjugated Streptavidin
Conjugate:	Cy5™
F/P Ratio:	10.0

Target Details

Purity/Specificity:	Cy5 conjugated Streptavidin was prepared from chromatographically purified streptavidin followed by conjugation to the dye Cy5 and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Streptavidin. No reaction was observed against anti-Avidin.
----------------------------	--

Application Details

Tested Applications:	Dot Blot, WB
Suggested Applications:	ELISA, EM, FISH, Functional Assay, IF, IHC, Microarray, Multiplex (Based on references)
Application Note:	Streptavidin Cy5 Conjugated has been tested by dot blot and western blot and is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.
Assay Dilutions:	All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.
FC:	1:500 - 1:2,500
FLISA:	1:10,000 - 1:50,000
IF:	1:1,000 - 1:5,000

Formulation

Physical State:	Lyophilized
Concentration:	1.0 mg/mL by UV absorbance at 280 nm
Buffer:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Preservative:	0.01% (w/v) Sodium Azide
Stabilizer:	10 mg/mL Bovine Serum Albumin (BSA) - Immunoglobulin and Protease free
Reconstitution Volume:	1.0 mL
Reconstitution Buffer:	Restore with deionized water (or equivalent)

Shipping & Handling

Shipping Condition:	Ambient
Storage Condition:	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Expiration:	Expiration date is one (1) year from date of receipt.


Images



Bottle


Streptavidin Cy5 Conjugated

A Genome Imaging




Fluorescence in situ Hybridization (FISH)

Genome imaging and chromosome differences
 (A) Nucleus of a primary human fibroblast imaged using 3D FISH with the maternal and paternal copies of Chromosome 6, 8, and 11 painted red, green, and white, respectively (left). Subsection highlighting the separation between the maternal and paternal copies of Chromosome 11, now colored red (right). Streptavidin Cy5 Conjugated (p/n S000-06). Fig 2. PMID: 34877507.



Immunofluorescence Microscopy

Immunohistochemical staining of synaptic proteins PSD-95 and synaptophysin. Panel a shows representative immunohistochemical staining of dorsolateral striatal tissues from all four groups with an antibody against PSD-95 protein (red) and the nuclear stain DAPI (blue). The insert boxes in each panel are images at a higher magnification showing distinct puncta (synaptic staining). The scales bars in each set of panels represent 10 microns. Panel b shows representative immunohistochemical staining of dorsolateral striatal tissues from all four groups with an antibody against synaptophysin protein (red) and the nuclear stain DAPI (blue). Antibodies used to detect PSD-95 and synaptophysin and biotinylated goat anti-mouse IgG. Sections incubated with Cy5-conjugated streptavidin (p/n S000-06) and Alexa Fluor 594-conjugated goat anti-rabbit IgG. Fig 5. PMID: 24316165.



Western Blot

Western Blot of Cy5 conjugated Streptavidin. Lane 1: Biotin conjugated Guinea Pig IgG. Lane 2: none. Load: 100 ng per lane. Primary antibody: none. Secondary antibody: Cy5 conjugated Streptavidin at 1:1,000 for 60 min at RT. Block: MB-070 for 30 min at RT. Predicted/Observed size: 55 kDa, 28 kDa for Guinea Pig IgG. Other band(s): none.

References

- Reed KSM et al. Genome reorganization and its functional impact during breast cancer progression. *bioRxiv [preprint]* (2025)
- Lindsly, S et al. Functional organization of the maternal and paternal human 4D Nucleome. *IScience* (2021)
- Seitter H. et al. Voltage-gated calcium channel subunit $\alpha 2\delta$ -3 shapes light responses of mouse retinal ganglion cells mainly in low and moderate light levels. *bioRxiv* (2020)
- Li H et al. Snap chip for cross-reactivity-free and spotter-free multiplexed sandwich immunoassays. *J Vis Exp.* (2017)
- Zhu D et al. Novel glypican-3-binding peptide for in vivo hepatocellular carcinoma fluorescent imaging. *Bioconjug Chem.* (2016)
- Tikidji-Hamburyan A et al. Retinal output changes qualitatively with every change in ambient illuminance. *Nat Neurosci.* (2015)
- Hovhannisyan A et al. Effects of the jimpy mutation on mouse retinal structure and function. *J Comp Neurol.* (2015)
- Toy, WA et al. Treadmill exercise reverses dendritic spine loss in direct and indirect striatal medium spiny neurons in the 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) mouse model of Parkinson's disease. *Neurobiology of Disease* (2014)
- Gdula, MR et al. Remodeling of three-dimensional organization of the nucleus during terminal keratinocyte differentiation in the epidermis. *The Journal of Investigative Dermatology* (2013)
- Pla-Roca, M et al. Antibody colocalization microarray: a scalable technology for multiplex protein analysis in complex samples. *Molecular & Cellular Proteomics : Mcp* (2012)
- Fessing, MY et al. p63 regulates Satb1 to control tissue-specific chromatin remodeling during development of the epidermis. *The Journal of Cell Biology* (2011)

Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.