

Datasheet for WM9-01-0001**WM9 Viable Cells****Overview**

Description:	WM9 Viable Cells - WM9-01-0001
Item No.:	WM9-01-0001
Size:	1 million cells
Applications:	Cellular Assay, IF, IHC, Multiplex, Other, WB
Origin:	Human

Product Details

Background:	WM9 is a human metastatic melanoma cell line that was established from a metastatic site (left axillary node) in a male patient. This cell line features the V600E (Val600Glu) mutation at codon 600 in the BRAF gene. This mutation causes constitutively active kinase activity and activation of MEK and ERK signaling pathway. This cell line also expresses PTEN hemizygous deletion, and is wild type for N-RAS, c-KIT, and CDK4. WM9 cells produce xenograft tumors when injected into immunocompromised mice.
Synonyms:	Melanoma, patient derived tumor, tumor models, skin cancer, xenograft
Species of Origin:	Human

Target Details

Purity/Specificity:	Cells are sterile, validated by short tandem repeat profiling, and are tested as negative for mycoplasma. It is recommended that cell lines are tested for mycoplasma contamination and short tandem repeat (STR) profiling every 10 passages or each time a frozen seed stock is made. See cell culture protocol for additional details.
Relevant Links:	• Cell Line EULA • Melanoma Cell Culture Protocol

Application Details

Suggested Applications:	Cellular Assay, IF, IHC, Multiplex, Other, WB (Based on references)
--------------------------------	---

Application Note: The key applications of these cell lines include genetic studies, xenograft production, drug testing, and drug target discovery. These cell line models can be used in various biological assays, and for identifying critical target genes, and cell signaling pathways.

Assay Dilutions: All assays should be optimized by the user. Recommended dilutions (if any) may be listed below.

Cell Line Data

Cell Line:	Human Melanoma
Product Type:	Viable Cells
Cell Viability:	Yes
Stage:	Metastasis
BRAF:	V600E
CDK4:	WT
C-Kit:	WT
N-RAS:	WT
PTEN:	Hemizygous Deletion
Paired:	No
Medium:	Tumor Specialized Media with 2% HI-FBS
Sub-culture:	Cells should be maintained between 30 – 95% confluence in tumor specialized medium with 2% FBS; split cultures 1:5 every 7-14 days using 0.25% trypsin/EDTA.
Incubation:	36°C with 5% CO ₂

Formulation

Physical State:	Frozen Cell Suspension
Concentration:	1.0 million cells/mL Count By Hemocytometer
Buffer:	None
Preservative:	None
Stabilizer:	None

Shipping & Handling

Shipping Condition: Dry Ice

Storage Condition: Cells are frozen with 90% FBS/10% DMSO solution at about 1×10^6 cells/ml. Store vial in liquid nitrogen upon arrival.

Expiration: Expiration date is two (2) years from date of receipt.

Images



Flask

Human melanoma tumor cells with known gene mutations, disease stage, STR, and RPPA profiling

References

- Simiczjew A et al. Comparative Proteomic Analysis of the Secretome of Control and BRAF/MEK Inhibitor-Resistant Melanoma Cells. *J Proteome Res.* (2026)
- Kujawa T et al. Comprehensive transcriptomic characterization of two melanoma cell lines with acquired dual resistance to BRAF and MEK inhibitors. *Sci Rep.* (2026)
- Bauer JA et al. Nitrosylcobalamin Selectively Targets Tumors via Cobalamin Uptake and Lysosomal Processing. *Front Biosci (Elite Ed).* (2026)
- Wadzyska J et al. The impact of cellular elements of TME on melanoma biology and its sensitivity to EGFR and MET targeted therapy. *Biochim Biophys Acta Mol Cell Res.* (2023)
- Hippner M et al. Alpha-Enolase (ENO1) Correlates with Invasiveness of Cutaneous Melanoma—An In Vitro and a Clinical Study. *Diagnostics (Basel).* (2022)
- Kupczyk P et al. PARP1 as a Marker of an Aggressive Clinical Phenotype in Cutaneous Melanoma—A Clinical and an In Vitro Study. *Cells.* (2021)
- Malek N et al. The origin of the expressed retrotransposed gene ACTBL2 and its influence on human melanoma cells' motility and focal adhesion formation. *Sci Rep.* (2021)
- Makowiecka A et al. Thymosin β 4 regulates focal adhesion formation in human melanoma cells and affects their migration and invasion. *Front Cell Dev Biol.* (2019)
- Dratkiewicz E, et al. Characterization of Melanoma Cell Lines Resistant to Vemurafenib and Evaluation of Their Responsiveness to EGFR- and MET-Inhibitor Treatment. *Int J Mol Sci* (2019)
- Makowiecka A et al. Varying effects of EGF, HGF and TGF β on formation of invadopodia and invasiveness of melanoma cell lines of different origin. *Eur J Histochem.* (2016)

Disclaimer

No test method can provide total assurance that the hepatitis B virus, hepatitis C virus, human immunodeficiency virus, or any other infectious agents are absent. Thus, all blood products, including purified proteins derived from human blood sources, should be handled at Biosafety Level 2 as recommended by the CDC\NIH manual entitled Biosafety in Microbiological and Biomedical Laboratories for potentially infectious human serum, blood specimens or proteins derived from same. Source material for the human blood product supplied to your facility has been tested for the detection of HIV antibody, Hepatitis B surface antigen, antibody to Hepatitis C, HIV 1 antigen(s), antibody to HTLV - I/II, and syphilis by FDA guidelines. All units were found to be non-reactive/negative for these tests. All human blood source material is collected in FDA licensed centers and is tested with FDA approved test kits.

Cell Line Limited Use License Required. THIS PRODUCT IS SUBJECT TO AN END-USER LICENSE AGREEMENT (EULA). BY ACCEPTING THIS PRODUCT, RECIPIENT AGREES TO BE BOUND BY THE TERMS OF USE SET FORTH BELOW and SET FORTH IN THE EULA. THIS PRODUCT IS FOR IN VITRO RESEARCH USE ONLY. THERAPEUTIC, DIAGNOSTIC, OR VETERINARY USE IS PROHIBITED. This product may not be resold or transferred by the recipient and may be used only by the recipient, in the recipient's facility and only for research use and other uses specifically permitted by the EULA. No other commercial use is allowed. "Commercial Use" means any and all uses of this product by recipient or others for monetary or other consideration, including providing services, supplying information or data to unaffiliated third parties, and resale or transfer of this product for any use. Recipient has no right to modify, derivatize, genetically engineer or otherwise create variations of this product or associated cells or cell lines. ROCKLAND AND WISTAR MAKE NO REPRESENTATIONS AND EXTEND NO WARRANTIES OF ANY KIND, EITHER EXPRESSED OR IMPLIED, INCLUDING AS TO MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, OR THAT THE USE OF THE PRODUCTS WILL NOT INFRINGE ANY PATENT, COPYRIGHT, TRADEMARK, OR OTHER PROPRIETARY RIGHTS. The terms set forth herein and in the EULA shall be governed by the laws of the Commonwealth of Pennsylvania, USA. To obtain a COMMERCIAL USE license for this product, please contact Rockland Immunochemicals, Inc. Please contact a technical service representative for more information. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.